arXiv:1602.07841v1 [astro-ph.SR] 25 Feb 2016

Submitted to ApJ

The Distance Measurement of NGC 1313 with Cepheids

Gao Qing[1], Wei Wang[1], Ji-Feng Liu[1]

ABSTRACT

We present the detection of Cepheids in the barred spiral galaxy NGC 1313, using the Wide Field and Planetary Camera 2(WFPC2) on *Hubble Space Telescope*. Twenty B(F450W) and V(F555W) epochs of observations spanning over three weeks were obtained, on which the profile-fitting photometry of all stars in the monitored field was performed using the package HSTphot. A sample of 26 variable stars have been identified to be Cepheids, with periods between 3 and 15 days. Based on the derived period-luminosity (PL) relations in B- and V-bands, we **obtain** an extinction-corrected distance modulus of $\boldsymbol{\mu_{\rm NGC1313} = 28.32 \pm 0.10\,(\rm random) \pm 0.06\,(\rm systematic)}$, employing respectively the Large Magellanic Cloud (LMC) as distance zero point calibrators. The above moduli correspond to a distance of **4.61$\pm$0.21 (random)$\pm$0.13 (systematic) Mpc**, consistent with previous measurements reported in the literature within uncertainties. In addition, the reddening to NGC 1313 is found to be small.

Subject headings: Cepheids - distance scale - galaxies: individual (NGC1313, LMC)

1. INTRODUCTION

Cepheids are luminous variable stars with **radial** pulsations, which have been proven to be reliable distance indicators with the well-established period-luminosity relation in both observational and theoretical aspects. Given their intrinsic visual brightness range from -2 to -6 mag, Cepheids are ideal standard candles at Galactic and extragalactic distances, and provide useful calibrations for secondary extragalactic distance indicators including for example Tully-Fisher relation and planetary nebular luminosity functions.

[1]Key Laboratory of Optical Astronomy, National Astronomical Observatories, Chinese Academy of Sciences, Beijing 100012, China

Thanks to the improved resolution of the *Hubble Space Telescope (HST)*, many Cepheids have been discovered in the galaxies in the **Local Supercluster** and the distances to these galaxies have therefore been determined. This campaign extends the distance scale that can be determined reliably by a factor of 10 as compared to the pre-*HST* era, when Cepheids were mostly detected and studied from the ground. For a more detailed understanding of Cepheids distance scale and its implications to the local universe, the readers are referred to the classic review by Madore & Freedman (1991).

The nearby starburst galaxy NGC 1313 is a spiral galaxy with morphological type SB(s) (de Vaucouleurs et al. 1991). It is often described as a galaxy in transition between the Magellanic type galaxies and normal disk galaxies such as M33, given its high rotation velocity and very little or no radial metallicity gradient (Mollá & Roy 1999). Being very active in star formation, NGC 1313 and its super shell nebula have been extensively studied (e.g. Ryder 1995; Pellerin et al. 2007), as well as the X-ray sources NGC1313 X-1 and X-2 (Feng & Kaaret 2005, 2006; Liu et al. 2007, 2012) and typeII supernova SN1978K (Ryder et al. 1993). The distance to this galaxy has been determined previously using the methods including for example supernova radio flux density and H II region size distribution, which are listed in Table 1 and show modest diversity. Securing an independent distance measurement using Cepheids in NGC 1313 is necessary in this context.

2. Observation and data analysis

The sky region around NGC 1313 X-2 was monitored 20 times over three weeks in 2008 with *H*ST WFPC2 in the F450W and F555W filters for the program GO-11227 (PI: Jifeng Liu), which was aimed to search and identify periodicity of NGC 1313 X-2 in the optical. For each observation, two 500 s exposures in F450W were taken, followed by two shots of 400 s and 700 s in F555W. The observations were calibrated with the best calibration files on 2008 August 26, and standard photometric analysis were performed on each frame using the photometry package HSTphot (Dolphin 2000), which is designed to handle the undersampled point-spread functions (PSF) in WFPC2 images. The package conducts procedures for cleaning cosmic rays, removing hot pixels, fitting PSF, and applying aperture corrections to secure precise photometry in the VEGAmag photometric system.

In total, 32482 sources are detected and identified as well-photometered stars in the 5.7 arcmin^2 of NGC 1313 covered by the WFPC2 field-of-view. Further investigations will be based on them.

2.1. Variable Star Search and Cepheids Identification

Only well-photometered point sources are included for the search of variable stars. Following a similar approach as Dolphin et al. (2003) in their study of variables in Sextans, we define a variable star with magnitude variation $\sigma_{\rm mag} \geqslant 0.15$ mag and chi square $\chi^2_\nu \geqslant 2$, where

$$\sigma_{mag} \equiv (\frac{1}{40}(\sum_{i=1}^{20}(B_i - \overline{B})^2 + \sum_{i=1}^{20}(V_i - \overline{V})^2))^2$$

and

$$\chi^2_\nu \equiv (\frac{1}{40}(\sum_{i=1}^{20}\frac{(B_i - \overline{B})^2}{\sigma^2_{i,B}} + \sum_{i=1}^{20}\frac{(V_i - \overline{V})^2}{\sigma^2_{i,V}}))^{\frac{1}{2}}$$

Periods of these variable star candidates are determined using a modified Lafler-Kinman algorithm (Lafler & Kinman 1965), but with a relatively loose criterion of minimum $\Theta \leqslant 0.85$ in either B or V light curves for in order not to miss any periodic variables. Among the total 32, 482 stars, 1616 are found to be variables, and $\sim 65\%$ of them show periodic variations.

The light curves of these variable stars with periodic variations were visually examined, and 26 of them are found to be Cepheid on the basis of their distinctively rapid brightening, followed by a long, linear decay phase. They are presented in the color-magnitude diagram (CMD) as in Fig. 1 as red filled circles, along with all 32, 482 stars in black dots.

2.2. Light curves and average magnitudes for Cepheids

In Table 2 and 3, V- and B-bands time-series photometry for the Cepheids are presented for each epoch, along with photometric uncertainties, respectively. **We determined the pulsation period and the average magnitudes in V- and B- bands for each Cepheid by using template fitting procedures. The template fitting method based on Fourier analysis and Principal component analysis (PCA) was firstly introduced by Stetson (1996) and Kaluzny, J. and Stanek, K. Z. and Krocker (1998). It can obtained more robust periods and precisely measure average magnitudes even in few observational epochs and bad-sampled light curves. Yoachim, P. and McCommas, L. P. and Dalcanto (2009) and Pejcha, O. and Kochanek, C. S. (2012) expanded on these methods to build new templates for analysis of HST Cepheids data. The templates of Yoachim, P. and McCommas, L. P. and Dalca (2009) include BVI bands and short-period Cepheids ($<$ 10 days) in the analysis. It was ideal for characterizing our discovered Cepheids.**

The training set included the light curves of fundamental (FU) type Cepheids stars comes from OGLE database for the LMC (Udalski et al. 1999a), SMC (Udalski et al. 1999b) and VizieR database for the Milky Way (Ochsenbein, F. and Bauer, P. and Marcout, J. 2000). We put in a single fake I- band observation as there are no I band photometry. Considering there have fewer data points in B- band and deblending problems can cause outlier points in light curves, we fixed all additional PCA parameters in fitting procedure. As a result, the magnitudes and periods were determined through template fitting. The best-fit light curve templates also generated from obtained parameters.

The light curves of photometry data along with their best-fit templates (solid lines) in V- and B- are plotted in Figure 2 with their Ids, periods, V and B magnitudes. The average V- and B- magnitudes and their measurement uncertainties for each Cepheid are given in Table 4. The χ^2/dof for the fitting are given in the last column.

3. PL relations and the distance to NGC 1313

The phase-weighted apparent V and B PL relations for the Cepheids in NGC 1313 are plotted in Figures 3. The filled circles show the 24 Cepheids used for the distance calculating, while the 2 filled diamonds show the excluded ones, namely Cepheid 10 and 11. The NGC 1313 data are superimposed on the PL relation for LMC Cepheids (open circles) from updated OGLE II catalog (Udalski et al. 1999a). The solid lines are fits to the LMC data.

To determine the distance to the galaxy NGC 1313, the apparent PL relations need to be calibrated with the absolute PL relations in LMC galaxies, which contain a large number of Cepheids and have a well determined distance modulus of $18.493 \pm 0.008\,(\mathrm{statistical}) \pm 0.047\,(\mathrm{systematic})$ **(Pietrzyński et al. 2013). From the OGLE II LMC Cepheid catalog (Udalski et al. 1999a), we use their revised PL relations which using 650 FU Cepheids in V- band and 662 FU Cpeheids in B- band :**

$$\begin{aligned} M_V(\mathrm{LMC}) &= (-2.779 \pm 0.031)\log P + (17.066 \pm 0.021) \\ M_B(\mathrm{LMC}) &= (-2.439 \pm 0.046)\log P + (17.368 \pm 0.031) \end{aligned} \tag{1}$$

with **standard deviation of 0.160 mag at V and 0.239 mag at B**, respectively. **Here $M_V(\mathrm{LMC})$ and $M_B(\mathrm{LMC})$ are extinction-corrected values using the reddening of E($B-V$)=0.147 to LMC.**

In order to avoid incompleteness bias in fitting a slope to the NCG 1313 data, only the intercept of the linear regression was fitted, while keeping the slope fixed to the LMC values. We shift the NGC 1313 magnitudes to match those of LMC, and determine the relative distance moduli by minimizing the best-fit chi square of the linear fitting of the combined data set. As a result, the

apparent relative distance moduli between NGC 1313 and the LMC are measured to be $\boldsymbol{\Delta\mu_V = 9.893 \pm 0.049}$ mag and $\boldsymbol{\Delta\mu_B = 9.872 \pm 0.057}$ mag.

3.1. The true distance modulus and its uncertainties

The above derived distance moduli need to be corrected for foreground extinction and reddening effect. **The true absolute wavelength-independent distance modulus to NGC 1313 can be expressed as a function of the relative apparent modulus $\Delta\mu$ between NGC 1313 and LMC, the LMC distance modulus, and the extinction for NGC 1313, as:**

$$\boldsymbol{\mu_V = \Delta\mu_V + \mu^{\rm LMC} = \mu + A_V} \tag{2}$$

$$\boldsymbol{\mu_B = \Delta\mu_B + \mu^{\rm LMC} = \mu + A_B} \tag{3}$$

$$\boldsymbol{\mu = \mu_V - R_V(\mu_B - \mu_V)} \tag{4}$$

Where the reddening law of $R_V = A_V/E(B-V) = 3.3$ for NGC 1313 and the Magellanic Clouds (Ferrarese et al. 1996). The true modulus to $\boldsymbol{\mu^{LMC} = 18.493}$ **mag (Pietrzyński et al. 2013). By subtracting Equation 3 to Equation 2, we yield the extinction of NGC 1313:**

$$\boldsymbol{E(B-V) = A_B - A_V = \Delta\mu_B - \Delta\mu_V} \tag{5}$$

Based on Equation 3 and Equation 2, we found the apparent distance moduli for NGC 1313 are then $\mu_V = 28.386 \pm 0.049$ mag and $\mu_B = 28.365 \pm 0.057$ mag. Those two modulus agree to each other within errors. That implied zero reddening. We computed an error-weighted average value as the true distance modulus to NGC 1313 $\mu = 28.377 \pm 0.038$ mag instead of applying dereddening precedure of Equation 4.

In the above results,errors for the distance modulus consist of two independent factors: random errors and systematic errors. The random errors are calculated using Equation (18) of Saha et al. (2000), propagated mainly from the errors on stellar photometry and intrinsic widths of the PL relation in V and B:

$$\boldsymbol{\sigma^2_{random} = (\sigma^2_V + \sigma^2_B) + (\sigma^2_{\rm PL,B} + \sigma^2_{\rm PL,V})} \tag{6}$$

The first item in the right side is the estimated photometric errors for the mean V and B magnitudes, as propagated through the dereddening procedure, where $\boldsymbol{\sigma_V = 0.029}$ and $\boldsymbol{\sigma_B = 0.070}$ mag respectively. The second item is associated with the intrinsic width of the PL relation (i.e., a given Cepheid may not be on the mean ridge line of the PL relation), which is $\boldsymbol{\sigma_{\rm PL,V} = 0.046}$ mag, and $\boldsymbol{\sigma_{\rm PL,B} = 0.050}$ mag.

Several systematic errors also come into play: uncertainties due to the distance modulus of zero point galaxy, which is $0.008\ (\mathrm{statistical}) \pm 0.047\ (\mathrm{systematic})$ mag for LMC (Pietrzyński et al. 2013), errors derived from the observed scatter in PL relation of LMC and (all lower than 0.02 mag), uncertainties due the metallicity dependency of the PL relation. Walsh & Roy (1997) performed multifiber emission-line spectrophometry of 33 H II regions in NGC 1313, and derived a flat disc metallicity of $\log_{10}(\mathrm{O/H}) = -3.6 \pm 0.1$, which is very similar to that for LMC with $\log_{10}(\mathrm{O/H}) = -3.5 \pm 0.08$ **(Ferrarese et al. 2000)**. **On the observational side, in the joint fits of LMC, M81, and NGC 4258, Gerke, J. R. and Kochanek, C. S. and Prieto, J. L. and Stanek, K. Z. and (2011) found a luminosity dependence on metallicity of $\gamma_\mu = -0.62^{+0.31}_{-0.35}$ mag dex^{-1}. Applying this correction, our Cepheid distance moduli are corrected by: $\Delta\mu_{met} = (0.62 \pm 0.35)([O/H]_{NGC1313} - 8.5) = -0.062 \pm 0.035$. On the theoretical side, Bono, G. and Caputo, F. and Ma (2010) used the pulsation models with nebular oxygen abundance $7.56 \leq 12 + log(O/H) \leq 8.88$. It shows the metallicity effect on $\Delta\mu_0(WBV)$ is quite strong and they found $\gamma(WBV) = -0.58 \pm 0.03$ mag dex^{-1}. Applying this correction, our Cepheid distance moduli are corrected by: $\Delta\mu_{met} = (0.58 \pm 0.03)([O/H]_{NGC1313} - 8.5) = -0.058 \pm 0.003$.**

We list all our known sources of error in table 5. The distance modulus based on our 24 Cepheids is 28.32 ± 0.10 (random)± 0.06 (systematic), giving a corresponding distance of 4.61 ± 0.21 (random)± 0.13 (systematic) Mpc.

4. Discussion

We have derived a Cepheid distance to NGC 1313 based on V- and B-bands *H*ST WFPC2 observations. A total of 26 Cepheids have been found with periods ranging from **3.12 to 14 days**, with their phase-weighted average magnitudes in both bands and periods in V determined. Apparent V and B distance moduli to NGC 1313 have been derived, employing the LMC Cepheids samples to be zero point reference galaxies. After applying extinction corrections, the true distance modulus to NGC 1313 is **28.32 ± 0.10 (random)± 0.06 (systematic)** using LMC for the calibration of distance. These moduli correspond to distance of **4.61 ± 0.21 (random)± 0.13 (systematic) Mpc.** The Cepheid distance found in this work is slightly larger than previous measurements, but are consistent within errors with most results, especially those obtained using tip of RGB stars (summarized in Table 1).

An important factor on distance measurement is that Cepheids are oscillating in two modes — fundamental (FU) and first over-tone (FO). Cepheids with periods shorter than 8 days are likely to be oscillating as a first over-tone pulsator that appears brighter and bluer than fundamental Cepheids. To confuse this two type Cepheids would make the distance underestimated. Since the PL relations are calculated by only using FU Cepheids from OGLE database, we must make sure

that our 26 Cepheids are all belong to this type. Beaulieu et al. (1995) studied large sample of Cepheids light-curve in 490 nm in LMC, and found that the first over-tone (FO) Cepheids have a strong tendency to be more symmetric, and having lower amplitudes in light curves than the FU Cepheids. Fig. 4 presents the amplitudes against their periods of 26 Cepheids in V and 6 in B with detections in all 20 observations, along with the 94 LMC Cepheids from Beaulieu et al. (1995). It is obvious that their LMC FU and FO Cepheids occupy quite different regions in the Amplitude-Period (AP) plane. All of our Cepheids are within the area where LMC FU Cepheids are located, and they should be of FU type.

The extinction of NGC 1313 is also obtained when deriving the distance modulus. Based on **Equation 5**, we found the extinction to NGC 1313 is $\boldsymbol{E(B-V) = -0.01 \pm 0.07}$. The value for the estimated extinction is low compare with previous studies around the region of NGC 1313 X-2. An extinction of E(B-V) = 0.11 mag can be obtained from the Schlegel et al. (1998) dust map as the Galactic average value within 0.6° of NGC1313 X-2. An extinction of E(B-V) = 0.44 mag (corresponding to $n_H = 2.7 \times 10^{21} cm^{-2}$) is inferred from the power-law fit to the X-ray spectrum of NGC 1313 X-2 (Miller et al. 2003), which also includes contribution from circumbinary materials. The extinction value we obtained (< 0.03) through Cepheids suggests that the average extinction within $2'$ of NGC1313 X-2 is slightly smaller than the Galactic average value within 0.6° of NGC1313 X-2.

Acknowledgments

We thank Drs. Yu Bai, Yi Hu Song Wang and Hang Gong for very helpful discussions. JFL acknowledges the support from National Science Foundation of China through NSFC grants #11233004 and #11333004. WW is partially supported by NSFC grant #11203035 and 11233004 for this work.

REFERENCES

Beaulieu, J. P., et al. 1995, A&A, 303, 137

de Vaucouleurs, G., de Vaucouleurs, A., Corwin, Jr., H. G., Buta, R. J., Paturel, G., & Fouque, P. 1991, S&T, 82, 621

Dolphin, A. E. 2000, PASP, 112, 1383

Dolphin, A. E., et al. 2003, AJ, 125, 1261

Feng, H., & Kaaret, P. 2005, ApJ, 633, 1052

—. 2006, ApJ, 650, L75

Ferrarese, L., et al. 1996, ApJ, 464, 568

—. 2000, ApJS, 128, 431

Grisé, F., Pakull, M., Soria, R., Motch, C., Smith, I., Ryder, S., & Böttcher, M. 2008, arXiv preprint arXiv:0805.4011

Issa, I. A. 1981, Astronomische Nachrichten, 302, 251

Jacobs, B. A., Rizzi, L., Tully, R. B., Shaya, E. J., Makarov, D. I., & Makarova, L. 2009, AJ, 138, 332

Lafler, J., & Kinman, T. D. 1965, ApJS, 11, 216

Liu, J., Orosz, J., & Bregman, J. N. 2012, ApJ, 745, 89

Liu, J.-F., Bregman, J., Miller, J., & Kaaret, P. 2007, ApJ, 661, 165

Madore, B. F., & Freedman, W. L. 1991, PASP, 103, 933

Miller, J. M., Fabbiano, G., Miller, M. C., & Fabian, A. C. 2003, ApJ, 585, L37

Mollá, M., & Roy, J.-R. 1999, ApJ, 514, 781

Pellerin, A., Meyer, M., Harris, J., & Calzetti, D. 2007, ApJ, 658, L87

Pietrzyński, G., et al. 2013, Nature, 495, 76

Rizzi, L., Tully, R. B., Makarov, D., Makarova, L., Dolphin, A. E., Sakai, S., & Shaya, E. J. 2007, ApJ, 661, 815

Ryder, S., Staveley-Smith, L., Dopita, M., Petre, R., Colbert, E., Malin, D., & Schlegel, E. 1993, ApJ, 416, 167

Ryder, S. D. 1995, ApJ, 444, 610

Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, ApJ, 500, 525

Stetson, P. B. 1996, PASP, 108, 851

Bono, G. and Caputo, F. and Marconi, M. and Musella, I. 2010, ApJ, 715, 277

Gerke, J. R. and Kochanek, C. S. and Prieto, J. L. and Stanek, K. Z. and Macri, L. M. 2011, ApJ, 743, 176

Kaluzny, J. and Stanek, K. Z. and Krockenberger, M. and Sasselov, D. D. and Tonry, J. L. and Mateo, M. 1998, AJ, 115, 1016

Ochsenbein, F. and Bauer, P. and Marcout, J. 2000, A&AS, 143, 23

Pejcha, O. and Kochanek, C. S. 2012, ApJ, 748, 107

Yoachim, P. and McCommas, L. P. and Dalcanton, J. J. and Williams, B. F. 2009, AJ, 137, 4697

Tully, R. B. 1988, Nearby galaxies catalog

Tully, R. B., et al. 2006, AJ, 132, 729

Udalski, A., Soszynski, I., Szymanski, M., Kubiak, M., Pietrzynski, G., Wozniak, P., & Zebrun, K. 1999a, Acta Astron., 49, 223

—. 1999b, Acta Astron., 49, 437

Walsh, J. R., & Roy, J.-R. 1997, MNRAS, 288, 726

Weiler, K. W., van Dyk, S. D., Montes, M. J., Panagia, N., & Sramek, R. A. 1998, ApJ, 500, 51

This preprint was prepared with the AAS LaTeX macros v5.2.

Table 1:: The previous and current determinations of the distance to NGC 1313 using various methods, as listed in Column 2. Only measurements uncertainties (random) are taken into account in the error budgets given in Column 1 except this work.

Distance (Mpc)	Method	References
$4.13^{+0.11}_{-0.12}$	Tip of RGB (TRGB)	Bryan Mendez et al 2002
$4.15^{+0.06}_{-0.06}$	TRGB	Tully et al. (2006)
$4.27^{+0.06}_{-0.06}$	TRGB	Rizzi et al. (2007)
$4.07^{+0.20}_{-0.21}$	TRGB	Grisé et al. (2008)
$4.39^{+0.04}_{-0.04}$	TRGB	Jacobs et al. (2009)
3.7[a]	Tully-Fisher correlation	Tully (1988)
3.0	HII region size distribution	Issa (1981)
$3.9^{+0.8}_{-0.8}$	SN 1978K radio flux density	Weiler et al. (1998)
$\mathbf{4.61 \pm 0.21 \pm 0.13}$	Cepheids, calibrated with LMC	this work

[a] Assuming the Hubble constant H_0=75.0 km s^{-1} Mpc^{-1}.

Table 2:: V photometry of the 26 Cepheids discovered in this work.

JD	C1	C2	C3	C4	C5	C6
54607.92485	24.489±0.073	24.841±0.074	24.352±0.063	25.072±0.101	24.726±0.130	24.450±0.049
54608.05749	24.276±0.065	24.871±0.074	24.250±0.053	24.957±0.072	24.585±0.108	24.354±0.048
54609.0561	24.007±0.054	24.450±0.054	24.461±0.053	24.649±0.070	24.807±0.092	24.420±0.047
54610.05472	24.566±0.092	24.468±0.055	24.686±0.064	24.517±0.054	24.854±0.091	23.983±0.037
54611.05263	24.579±0.062	24.536±0.059	24.918±0.087	24.719±0.061	25.076±0.108	23.814±0.035
54612.11791	24.728±0.109	24.468±0.077	25.206±0.093	24.663±0.059	25.559±0.180	23.958±0.044
54613.24291	24.874±0.121	24.755±0.086	25.025±0.145	24.857±0.082	24.872±0.089	24.173±0.047
54614.24152	24.577±0.078	24.969±0.103	24.259±0.054	24.806±0.078	24.591±0.071	24.084±0.040
54615.24083	24.092±0.062	24.915±0.091	24.709±0.082	24.931±0.070	24.864±0.101	24.309±0.045
54616.11097	24.468±0.079	24.707±0.066	24.826±0.080	25.067±0.094	24.903±0.103	24.287±0.044
54617.16791	24.600±0.071	24.295±0.050	24.947±0.093	24.734±0.063	25.062±0.166	24.354±0.047
54618.7061	24.787±0.106	24.595±0.062	25.252±0.111	24.394±0.049	25.504±0.161	24.487±0.053
54619.23597	24.877±0.101	24.599±0.062	24.688±0.083	24.583±0.067	24.967±0.096	24.668±0.061
54620.16444	24.974±0.118	24.728±0.066	24.258±0.052	24.602±0.060	24.704±0.098	24.561±0.057
54621.23388	24.375±0.060	24.897±0.080	24.830±0.084	24.834±0.068	24.945±0.112	24.607±0.065
54622.16235	24.266±0.057	25.306±0.141	24.944±0.076	24.803±0.063	25.016±0.118	24.410±0.047
54623.16097	24.361±0.061	24.705±0.070	25.081±0.099	25.017±0.074	25.076±0.113	24.471±0.050
54624.08944	24.743±0.101	24.370±0.050	25.048±0.098	25.210±0.085	25.291±0.163	23.969±0.035
54625.15888	24.857±0.107	24.456±0.059	24.668±0.063	24.590±0.056	24.815±0.105	23.857±0.038
54626.62555	24.827±0.090	24.563±0.059	24.482±0.061	24.451±0.052	24.854±0.105	24.052±0.039

C7	C8	C9	C10	C11	C12	C13
24.566±0.054	23.678±0.044	23.803±0.033	24.760±0.088	24.518±0.063	24.988±0.119	25.464±0.149
24.540±0.053	23.601±0.034	23.799±0.043	24.581±0.104	24.582±0.077	24.899±0.089	25.450±0.168
24.268±0.044	23.681±0.044	23.640±0.030	24.424±0.065	24.763±0.076	24.641±0.070	25.415±0.128
24.202±0.044	23.709±0.046	23.824±0.040	24.373±0.056	24.526±0.073	24.406±0.059	24.751±0.074
24.360±0.050	23.839±0.040	23.973±0.041	24.688±0.077	24.370±0.060	24.673±0.077	25.412±0.172
24.461±0.052	23.876±0.041	24.084±0.042	...	24.543±0.098	24.826±0.085	25.177±0.111
24.484±0.052	24.050±0.047	24.213±0.054	24.344±0.057	24.952±0.126	24.870±0.090	25.543±0.160
24.633±0.059	24.131±0.048	24.371±0.049	24.399±0.059	24.495±0.064	24.997±0.098	25.376±0.139
24.419±0.049	24.210±0.061	24.458±0.063	24.795±0.095	24.515±0.066	24.108±0.079	24.927±0.090
24.160±0.041	24.065±0.048	24.395±0.053	24.701±0.072	24.568±0.070	24.522±0.064	25.078±0.164
24.331±0.048	23.745±0.038	24.303±0.060	24.483±0.066	24.694±0.079	24.533±0.073	25.238±0.124
24.508±0.055	23.632±0.036	24.030±0.049	24.461±0.065	24.467±0.066	24.784±0.124	25.567±0.188
24.481±0.054	23.683±0.037	23.869±0.037	24.757±0.083	24.752±0.084	25.046±0.109	...
24.578±0.054	23.637±0.034	23.747±0.033	24.597±0.067	24.686±0.073	24.464±0.075	25.022±0.109
24.443±0.053	23.783±0.037	23.799±0.034	24.306±0.056	24.680±0.073	24.248±0.053	25.020±0.094
24.079±0.039	23.843±0.075	24.053±0.052	24.440±0.058	24.521±0.089	24.552±0.066	25.224±0.122
24.284±0.045	23.951±0.045	24.087±0.042	24.593±0.069	24.715±0.077	24.700±0.080	25.323±0.143
24.308±0.045	24.136±0.048	24.130±0.048	24.760±0.075	24.799±0.080	24.889±0.086	25.360±0.126
24.413±0.050	24.171±0.051	24.358±0.049	24.325±0.056	24.495±0.067	24.890±0.105	25.110±0.106
24.566±0.061	24.170±0.052	24.409±0.072	24.528±0.067	24.668±0.077	24.227±0.054	25.124±0.135

Table 2:: V photometry of 26 Cepheids (continued)

JD	C14	C15	C16	C17	C18	C19
54607.92485	25.401±0.126	25.411±0.136	25.334±0.109	25.464±0.149	24.967±0.097	25.259±0.122
54608.05749	...	25.407±0.122	25.220±0.139	25.450±0.168	25.211±0.105	25.179±0.095
54609.0561	...	25.223±0.181	24.818±0.072	25.415±0.128	...	25.216±0.108
54610.05472	25.083±0.102	24.798±0.072	25.177±0.114	24.751±0.074	25.615±0.156	24.917±0.086
54611.05263	25.328±0.126	25.394±0.126	25.306±0.116	25.412±0.172	25.184±0.148	25.158±0.105
54612.11791	25.408±0.133	25.315±0.115	25.367±0.122	25.177±0.111	25.280±0.116	25.370±0.107
54613.24291	24.649±0.079	24.779±0.095	24.927±0.084	25.543±0.160	25.536±0.152	25.302±0.124
54614.24152	24.916±0.084	25.206±0.104	24.904±0.097	25.376±0.139	25.850±0.189	24.943±0.083
54615.24083	25.284±0.121	25.499±0.135	25.173±0.102	24.927±0.090	24.925±0.091	25.074±0.117
54616.11097	...	25.120±0.094	25.134±0.095	25.078±0.164	25.142±0.099	25.360±0.122
54617.16791	24.782±0.081	25.040±0.096	25.452±0.166	25.238±0.124	25.581±0.162	25.449±0.150
54618.7061	25.207±0.120	25.368±0.131	24.942±0.090	25.567±0.188	25.169±0.112	25.031±0.097
54619.23597	25.493±0.162	...	24.954±0.094	...	24.862±0.087	25.080±0.090
54620.16444	25.537±0.146	24.654±0.066	25.295±0.130	25.022±0.109	25.151±0.100	25.255±0.112
54621.23388	24.755±0.074	25.318±0.132	25.375±0.130	25.020±0.094	25.653±0.203	25.514±0.130
54622.16235	25.115±0.100	25.517±0.152	25.271±0.111	25.224±0.122	...	24.564±0.080
54623.16097	25.270±0.119	25.488±0.154	24.941±0.084	25.323±0.143	24.949±0.089	25.255±0.107
54624.08944	25.395±0.130	24.698±0.067	24.939±0.081	25.360±0.126	25.184±0.129	25.221±0.090
54625.15888	24.918±0.099	25.201±0.106	25.298±0.114	25.110±0.106	25.713±0.173	25.317±0.139
54626.62555	25.303±0.127	25.556±0.147	25.501±0.141	25.124±0.135	24.897±0.086	24.804±0.081

C20	C21	C22	C23	C24	C25	C26
25.390±0.170	25.141±0.091	25.551±0.113	25.153±0.086	25.315±0.102	25.789±0.151	25.830±0.193
25.152±0.117	25.430±0.119	25.386±0.130	25.142±0.080	25.427±0.115	25.824±0.154	25.702±0.142
25.640±0.177	25.609±0.154	25.983±0.163	25.250±0.093	25.334±0.112	24.959±0.078	25.225±0.087
25.673±0.171	25.766±0.179	25.740±0.134	25.217±0.084	25.076±0.084	25.194±0.091	25.339±0.102
25.366±0.151	25.376±0.120	25.381±0.108	24.564±0.055	25.109±0.104	25.570±0.142	25.693±0.176
25.415±0.217	25.358±0.164	25.614±0.151	24.795±0.065	25.289±0.114	25.586±0.133	...
25.537±0.162	25.710±0.163	25.873±0.159	25.089±0.078	25.205±0.096	25.064±0.084	25.115±0.082
25.187±0.112	25.937±0.191	25.563±0.137	25.457±0.126	25.493±0.122	25.247±0.099	25.667±0.126
25.688±0.193	25.129±0.095	25.810±0.150	25.448±0.103	25.012±0.091	25.410±0.109	25.482±0.154
25.873±0.207	25.596±0.137	26.054±0.179	24.520±0.052	25.170±0.092	...	25.912±0.160
25.149±0.114	25.656±0.159	25.174±0.094	24.632±0.062	25.412±0.129	25.245±0.138	25.014±0.083
25.427±0.178	25.791±0.201	25.993±0.189	25.150±0.087	...	25.087±0.090	25.858±0.155
...	25.272±0.115	25.740±0.154	25.270±0.098	24.789±0.072	25.399±0.130	25.835±0.154
24.827±0.081	25.493±0.126	25.141±0.085	25.466±0.103	24.958±0.078	25.504±0.118	25.953±0.158
25.554±0.153	25.739±0.155	25.417±0.105	24.601±0.055	25.116±0.087	25.601±0.128	25.288±0.093
...	25.861±0.173	25.718±0.133	24.675±0.057	25.185±0.092	25.106±0.093	25.468±0.106
24.821±0.088	25.140±0.095	25.435±0.109	25.097±0.085	25.523±0.135	25.234±0.096	25.699±0.134
25.555±0.206	25.511±0.126	25.382±0.105	25.108±0.080	24.834±0.075	25.490±0.113	26.039±0.167
25.698±0.185	25.812±0.174	25.919±0.166	25.333±0.095	25.238±0.099	25.770±0.157	25.105±0.082
25.068±0.106	25.655±0.168	25.183±0.104	24.604±0.057	25.429±0.118	25.056±0.086	25.498±0.113

Table 3:: B photometry of the 26 Cepheids discovered in this work.

JD	C1	C2	C3	C4	C5	C6
54607.92485	...	...	24.851±0.104	...	24.818±0.150	25.176±0.118
54608.05749	24.853±0.166	...	24.583±0.086	25.830±0.188	24.704±0.133	25.325±0.109
54609.0561	...	25.159±0.164	24.992±0.120	...	...	25.397±0.112
54610.05472	25.382±0.204	24.985±0.130	25.346±0.152	25.021±0.110	...	24.624±0.066
54611.05263	...	25.066±0.129	...	25.267±0.147	25.096±0.208	24.513±0.068
54612.11791	...	25.422±0.174	...	25.349±0.125	...	24.720±0.072
54613.24291	...	25.554±0.205	...	25.666±0.168	...	24.845±0.096
54614.24152	...	...	24.728±0.100	25.454±0.213	24.824±0.178	24.740±0.074
54615.24083	24.659±0.109	...	24.887±0.117	...	...	25.004±0.131
54616.11097	25.069±0.212	25.336±0.165	...	25.728±0.187	...	25.180±0.097
54617.16791	25.178±0.207	24.964±0.153	...	25.600±0.165	...	25.084±0.098
54618.7061	...	25.299±0.161	...	25.105±0.106	...	25.535±0.140
54619.23597	24.934±0.186	...	25.515±0.189	25.203±0.121	24.809±0.194	25.411±0.142
54620.16444	...	...	24.687±0.102	25.365±0.160	24.733±0.151	25.549±0.138
54621.23388	24.709±0.118	25.255±0.202	25.133±0.155	25.212±0.119	...	25.322±0.134
54622.16235	24.906±0.181	25.439±0.167	25.517±0.176	...	...	25.040±0.092
54623.16097	25.281±0.160	24.950±0.157	...	25.454±0.172	...	25.761±0.197
54624.08944	25.182±0.175	25.008±0.153	...	25.836±0.204	...	24.700±0.080
54625.15888	...	25.097±0.145	25.406±0.160	25.122±0.157	...	24.453±0.075
54626.62555	...	25.278±0.160	24.901±0.115	25.148±0.110	...	24.741±0.099

C7	C8	C9	C10	C11	C12	C13
25.095±0.129	24.326±0.085	24.434±0.057	24.997±0.153	25.009±0.131	25.360±0.196	25.422±0.156
24.764±0.084	24.165±0.066	24.399±0.058	25.101±0.148	25.199±0.172	25.307±0.192	25.506±0.147
24.384±0.104	24.333±0.101	24.230±0.060	24.701±0.103	24.998±0.129	25.063±0.144	...
24.529±0.091	24.329±0.074	24.662±0.069	24.708±0.102	...	24.807±0.152	...
24.813±0.105	24.560±0.091	24.782±0.077	25.017±0.157	25.019±0.141	25.032±0.151	25.991±0.199
24.713±0.083	24.902±0.129	25.092±0.101	...	...	25.359±0.210	25.347±0.131
24.966±0.115	24.866±0.155	25.034±0.092	24.937±0.172	...	...	25.865±0.203
24.953±0.101	24.947±0.140	25.333±0.147	24.551±0.134	24.987±0.192	...	25.996±0.203
24.778±0.087	24.989±0.139	25.243±0.126	...	25.051±0.163	24.679±0.108	25.834±0.181
24.401±0.065	24.623±0.142	25.054±0.090	...	25.305±0.183	24.832±0.154	25.056±0.097
24.835±0.095	24.547±0.105	25.197±0.110	24.679±0.147	...	24.792±0.176	25.916±0.204
24.757±0.085	24.219±0.067	24.868±0.094	24.823±0.160	25.101±0.186	25.221±0.180	...
24.927±0.132	24.399±0.096	24.539±0.071	25.074±0.199	25.021±0.149	...	...
24.911±0.097	24.283±0.071	24.219±0.053	25.334±0.193	...	24.943±0.135	25.108±0.098
24.875±0.094	24.585±0.090	24.540±0.065	24.869±0.149	25.063±0.141	24.895±0.127	25.633±0.145
24.435±0.068	24.612±0.148	24.773±0.085	24.731±0.116	25.084±0.145	...	...
24.628±0.078	24.703±0.103	24.838±0.087	25.123±0.195	...	25.350±0.208	25.780±0.170
24.685±0.080	25.159±0.160	24.873±0.083	25.218±0.176	...	25.385±0.214	24.985±0.090
25.016±0.106	25.033±0.144	25.175±0.106	24.819±0.163	...	25.160±0.173	25.535±0.143
24.917±0.098	24.818±0.115	25.370±0.116	24.664±0.100	24.737±0.115	24.753±0.128	...

Table 3:: B photometry of the 26 Cepheids discovered in this work.

JD	C14	C15	C16	C17	C18	C19
54607.92485	...	...	25.780±0.215	...	...	25.798±0.179
54608.05749	...	...	25.478±0.167	...	25.567±0.209	25.859±0.194
54609.0561	...	25.578±0.201	25.456±0.159	...	...	...
54610.05472	...	25.262±0.141	...	25.269±0.167	...	25.175±0.107
54611.05263	...	...	...	...	25.357±0.179	25.555±0.154
54612.11791	...	25.561±0.204	...	...	...	...
54613.24291	25.256±0.210	25.046±0.126	25.373±0.157	...	...	...
54614.24152	...	...	25.408±0.164	...	...	25.398±0.192
54615.24083	...	...	...	25.043±0.207	25.168±0.156	25.727±0.195
54616.11097	...	...	...	...	25.448±0.202	25.742±0.178
54617.16791	...	...	...	...	...	...
54618.7061	...	...	25.493±0.189	...	...	...
54619.23597	...	...	...	...	...	25.594±0.180
54620.16444	...	25.117±0.132	...	25.127±0.192	...	...
54621.23388	25.129±0.143	25.500±0.190	...	...	...	...
54622.16235	...	...	...	...	...	25.271±0.118
54623.16097	...	...	...	...	25.146±0.152	25.714±0.177
54624.08944	...	25.186±0.159	...	...	...	25.913±0.212
54625.15888	25.016±0.136	...	25.650±0.210	...	...	25.807±0.203
54626.62555	...	...	...	24.986±0.211	25.226±0.151	25.447±0.197

C20	C21	C22	C23	C24	C25	C26
...	25.632±0.188	...	25.678±0.213	...	...	...
...	...	...	25.708±0.176	...	...	...
...	...	...	25.656±0.152	...	25.437±0.147	...
...	...	...	25.790±0.162	25.611±0.186	...	...
25.167±0.167	25.528±0.186	...	24.915±0.090	25.343±0.148	...	...
...	25.467±0.201	...	25.516±0.134	...	...	...
...	...	...	25.714±0.161	...	25.276±0.169	25.455±0.172
...	...	...	...	...	25.479±0.163	...
...	...	...	25.720±0.185	25.184±0.134	...	...
...	...	...	24.931±0.089	25.563±0.182	...	...
...	...	...	25.253±0.115	...	...	25.667±0.178
...	...	...	25.901±0.184	...	...	...
...	...	...	25.817±0.188	...	...	...
...	...	25.722±0.166	25.804±0.182	25.512±0.197	...	...
...	...	25.676±0.168	25.053±0.096	...	...	25.594±0.165
...	...	...	25.053±0.096	...	25.190±0.124	...
25.046±0.148	...	...	25.426±0.130	...	...	...
...	25.584±0.197	...	...	...	...	...
...	...	...	...	...	...	25.391±0.138
25.325±0.195	...	25.554±0.146	24.953±0.089	...	25.497±0.161	...

Table 4:: The final sample of 26 Cepheids, with their IDs, coordinates and periods presented in Columns 1-4. Final photometry, photometric error, and number of observations in V are given in Columns 5-7, while those in B in Columns 8-10. **The χ^2/dof for the fitting are given in the last column.**

ID	R.A. (J2000)	Decl. (J2000)	P (days)	V (mag)	error (mag)	N_V	B (mag)	error (mag)	N_B	χ^2/dof
C1	3:17:57.12	−66:34:19.2	**6.50**	**24.540**	**0.029**	20	**25.172**	**0.088**	10	**2.6**
C2	3:18:23.76	−66:34:33.6	**7.39**	**24.663**	**0.018**	20	**25.326**	**0.051**	14	**1.4**
C3	3:18:23.76	−66:35:49.2	**5.86**	**24.721**	**0.031**	20	**25.246**	**0.071**	12	**3.0**
C4	3:18:30.96	−66:34:55.2	**8.00**	**24.747**	**0.020**	20	**25.446**	**0.048**	16	**1.7**
C5	3:18:20.16	−66:34:51.6	**5.85**	**24.970**	**0.023**	20	**25.180**	**0.076**	6	**0.8**
C6	3:18:28.8	−66:34:26.4	**14.00**	**24.297**	**0.019**	20	**25.133**	**0.040**	20	**3.3**
C7	3:18:25.92	−66:34:44.4	**6.54**	**24.405**	**0.021**	20	**24.777**	**0.038**	20	**3.4**
C8	3:18:20.64	−66:35:09.6	**10.50**	**23.894**	**0.012**	20	**24.644**	**0.027**	20	**1.5**
C9	3:18:26.4	−66:34:04.8	**11.50**	**24.098**	**0.013**	20	**24.892**	**0.023**	20	**1.7**
C10	3:18:18.72	−66:34:40.8	**4.00**	**24.541**	**0.022**	19	**24.909**	**0.047**	17	**1.8**
C11	3:18:18.96	−66:34:44.4	**3.75**	**24.601**	**0.030**	20	**25.057**	**0.079**	12	**3.3**
C12	3:18:18.96	−66:34:51.6	**5.50**	**24.629**	**0.021**	20	**25.119**	**0.048**	16	**1.5**
C13	3:18:26.16	−66:34:12.0	**4.08**	**24.841**	**0.022**	20	**25.668**	**0.043**	14	**1.2**
C14	3:18:22.32	−66:34:58.8	**3.90**	**25.150**	**0.036**	17	**25.669**	**0.107**	3	**0.7**
C15	3:18:25.44	−66:35:02.4	**3.56**	**25.130**	**0.029**	19	**25.575**	**0.071**	7	**1.3**
C16	3:18:26.16	−66:35:06.0	**5.00**	**25.175**	**0.022**	20	**25.796**	**0.065**	7	**0.9**
C17	3:18:22.08	−66:35:06.0	**5.24**	**25.215**	**0.025**	19	**25.587**	**0.087**	4	**0.8**
C18	3:18:22.32	−66:34:44.4	**3.79**	**25.314**	**0.027**	18	**25.716**	**0.068**	6	**0.9**
C19	3:18:26.4	−66:34:26.4	**4.14**	**25.126**	**0.027**	20	**25.648**	**0.054**	13	**1.0**
C20	3:18:22.8	−66:35:16.8	**2.99**	**25.416**	**0.052**	18	**25.641**	**0.134**	3	**1.4**
C21	3:18:24.0	−66:34:40.8	**3.93**	**25.571**	**0.024**	20	**25.816**	**0.068**	4	**0.4**
C22	3:18:26.4	−66:34:19.2	**3.12**	**25.630**	**0.040**	20	**26.038**	**0.130**	3	**1.9**
C23	3:18:30.72	−66:34:51.6	**5.00**	**24.965**	**0.025**	20	**25.474**	**0.045**	17	**2.0**
C24	3:18:23.76	−66:45:32.4	**4.50**	**25.191**	**0.028**	19	**25.732**	**0.090**	5	**1.4**
C25	3:18:28.32	−66:35:09.6	**4.29**	**25.360**	**0.021**	19	**25.851**	**0.057**	5	**0.7**
C26	3:18:27.36	−66:34:48.0	**3.99**	**25.551**	**0.046**	19	**26.127**	**0.134**	4	**1.0**

Table 5:: **Error Budget**

Source	Random Error	Systematic Error
Stellar Photometry	**0.076**	**...**
Widths of PL relation	**0.068**	**...**
μ_{LMC}	**0.008**	**0.047**
LMC PL relation	**...**	**0.011**
Metallicity Correction	**...**	**0.035**



Fig. 1.—: **Color-magnitude diagram for the HST/WFPC2 field under current study. Black dots represent all the detected stars, while the red filled circles represent the 26 Cepheids identified in this paper. The dashed lines represent the zero reddening instabillity strip of LMC Cepheids and its 2 σ width.**


C1
Per = 6.50 days <V> = 24.54 ± 0.03 mag
<B> = 25.17 ± 0.09 mag
red chi = 2.64
C2
Per = 7.39 days <V> = 24.66 ± 0.02 mag
<B> = 25.33 ± 0.05 mag
red chi = 1.46
C3
Per = 5.86 days <V> = 24.72 ± 0.03 mag
<B> = 25.25 ± 0.07 mag
red chi = 3.08
C4
Per = 8.00 days <V> = 24.75 ± 0.02 mag
<B> = 25.45 ± 0.05 mag
red chi = 1.78
C5
Per = 5.85 days <V> = 24.97 ± 0.02 mag
<B> = 25.18 ± 0.08 mag
red chi = 0.85
C6
Per = 14.00 days <V> = 24.30 ± 0.02 mag
<B> = 25.13 ± 0.04 mag
red chi = 3.36
C7
Per = 6.54 days <V> = 24.41 ± 0.02 mag
<B> = 24.78 ± 0.04 mag
red chi = 3.49
C8
Per = 10.50 days <V> = 23.89 ± 0.01 mag
<B> = 24.64 ± 0.03 mag
red chi = 1.54
< mag >
jd(days)
jd(folded)

Fig. 2.—: Light curves of the 26 Cepheids


C9
Per = 11.50 days <V> = 24.10 ± 0.01 mag
<B> = 24.89 ± 0.02 mag
red chi = 1.78
< mag >
jd(days)
jd(folded)


C10
Per = 4.00 days <V> = 24.54 ± 0.02 mag
<B> = 24.91 ± 0.05 mag
red chi = 1.88
< mag >
jd(days)
jd(folded)


C11
Per = 3.75 days <V> = 24.60 ± 0.03 mag
<B> = 25.06 ± 0.08 mag
red chi = 3.36
< mag >
jd(days)
jd(folded)


C12
Per = 5.50 days <V> = 24.63 ± 0.02 mag
<B> = 25.12 ± 0.05 mag
red chi = 1.50
< mag >
jd(days)
jd(folded)


C13
Per = 4.08 days <V> = 24.84 ± 0.02 mag
<B> = 25.67 ± 0.04 mag
red chi = 1.21
< mag >
jd(days)
jd(folded)


C14
Per = 3.90 days <V> = 25.15 ± 0.04 mag
<B> = 25.67 ± 0.11 mag
red chi = 0.75
< mag >
jd(days)
jd(folded)


C15
Per = 3.56 days <V> = 25.13 ± 0.03 mag
<B> = 25.58 ± 0.07 mag
red chi = 1.39
< mag >
jd(days)
jd(folded)


C16
Per = 5.00 days <V> = 25.18 ± 0.02 mag
<B> = 25.80 ± 0.06 mag
red chi = 0.91
< mag >
jd(days)
jd(folded)


C17
Per = 5.24 days <V> = 25.22 ± 0.03 mag
<B> = 25.59 ± 0.09 mag
red chi = 0.81
< mag >
jd(days)
jd(folded)


C18
Per = 3.79 days <V> = 25.31 ± 0.03 mag
<B> = 25.72 ± 0.07 mag
red chi = 0.94
< mag >
jd(days)
jd(folded)


C19
Per = 4.14 days <V> = 25.13 ± 0.03 mag
<B> = 25.65 ± 0.05 mag
red chi = 1.05
< mag >
jd(days)
jd(folded)


C20
Per = 2.99 days <V> = 25.42 ± 0.05 mag
<B> = 25.64 ± 0.13 mag
red chi = 1.48
< mag >
jd(days)
jd(folded)


C21
Per = 3.93 days <V> = 25.57 ± 0.02 mag
<B> = 25.82 ± 0.07 mag
red chi = 0.50
< mag >
jd(days)
jd(folded)


C22
Per = 3.12 days <V> = 25.63 ± 0.04 mag
<B> = 26.04 ± 0.13 mag
red chi = 1.98
< mag >
jd(days)
jd(folded)


C23
Per = 5.00 days <V> = 24.97 ± 0.02 mag
<B> = 25.47 ± 0.04 mag
red chi = 2.02
< mag >
jd(days)
jd(folded)


C24
Per = 4.50 days <V> = 25.19 ± 0.03 mag
<B> = 25.73 ± 0.09 mag
red chi = 1.48
< mag >
jd(days)
jd(folded)



Fig. 2.—: Light curves of the 26 Cepheids



FIG. 3a



FIG. 3b

Fig. 3.—: **(a) V- band PL relation for 26 NGC 1313 Cepheids (black dots), while the black filled diamonds show excluded ones. It plotted with respect to the LMC Cepheid (red open circles) shifted by 9.89 mag in relative apparent modulus. The magnitudes are plotted vs the logarithm of the period in units of days. The solid line is the shifted best fit to LMC Cepheid PL relation on V band. (b) B- band PL relation for 26 NGC 1313 Cepheids (black dots), while the black filled diamonds show excluded ones. It plotted with respect to the LMC Cepheid (red open circles) shifted by 9.87 mag in relative apparent modulus. The solid line is the shifted best fit to LMC Cepheid PL relation on B band.**



Fig. 4.—: Amplitude-period diagram for Cepheids including: 26 V-band NGC 1313 Cepheids (open circles), 6 B-band with detections **(filled circles)** in all 20 observations. NGC1313 Cepheids (filled dots), 52 LMC FO Cepheids (**filled** diamonds) and 32 LMC FU Cepheids (diagonal crosses) from Beaulieu et al. (1995). The NGC 1313 Cepheids clearly belong to the FU type.


C25
Per = 4.29 days <V> = 25.36 ± 0.02 mag
<B> = 25.85 ± 0.06 mag
red chi = 0.72
< mag >
jd(days)
jd(folded)


C26
Per = 3.99 days <V> = 25.55 ± 0.05 mag
<B> = 26.13 ± 0.13 mag
red chi = 1.01
< mag >
jd(days)
jd(folded)